



Long-Term Effect of Different Cropping Systems on Soil Physico-chemical Properties of Soil: Environmental Implications and Sustainability

Tejinder Kaur ^{a*}, K.B. Singh ^{b++} and Neeraj Rani ^c

^a Department of Soil Science, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

^b Punjab Agricultural Management and Extension Training Institute, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

^c School of Organic Farming, Punjab Agricultural University, Ludhiana, Punjab 141004, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author TK designed the study, performed the fieldwork, sampling, lab analysis, statistical analysis, wrote the protocol, and wrote the draft of the manuscript. Author KBS has contributed to study planning, manuscript conception and editing of the manuscript. Author NR has helped in the planning and execution of the experiment. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2025/v15i14687>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129861>

Original Research Article

Received: 11/11/2024

Accepted: 13/01/2025

Published: 18/01/2025

ABSTRACT

Rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) a dominant cropping system in Punjab Region, Northwestern India, overtime is believed to have negative effect on the soil and water quality. The present study was conducted in an ongoing long-term experiment at the research farm of the School of Organic Farming, Punjab Agricultural University, Ludhiana, to compare the effect of

⁺⁺Director;

*Corresponding author: E-mail: tejinderwaraich1234@gmail.com;

Cite as: Kaur, Tejinder, K.B. Singh, and Neeraj Rani. 2025. "Long-Term Effect of Different Cropping Systems on Soil Physico-Chemical Properties of Soil: Environmental Implications and Sustainability". *International Journal of Environment and Climate Change* 15 (1):225-41. <https://doi.org/10.9734/ijecc/2025/v15i14687>.

different cropping systems on soil physico-chemical properties. Ten treatments (cropping systems) were laid in a Randomized Complete Block Design (RCBD). The treatments were rice-wheat (CS₁), maize-wheat (CS₂), basmati rice-wheat-cowpea green manure (CS₃), maize-mustard-cowpea green manure (CS₄), maize-potato-spring groundnut (CS₅), maize-peas-spring groundnut (CS₆), maize+ cowpea (fodder)-maize fodder-oats fodder-sathi maize fodder (CS₇), sorghum multicut fodder-barseem fodder (CS₈), maize (cobs/fodder)-potato-onion (CS₉), and baby corn-potato-okra (CS₁₀). Soil samples were collected from four soil depths (0-7.5, 7.5-15, 15-22.5, and 22.5-30 cm) under each cropping system and analyzed for particles size, soil pH, EC, soil organic carbon, aggregate associated carbon, bulk density, porosity, water stable aggregates, MWD, and penetration resistance. Significantly lower soil pH was reported in CS₄ and CS₈ compared to other cropping systems. Soil EC was significantly higher in CS₅ and CS₁₀ while lowest in CS₄. Soil organic carbon (SOC), aggregate-associated carbon (AAC), and mean weight diameter (MWD) were significantly higher in CS₇ and CS₄. Soil bulk density (BD) and penetration resistance (PR) were significantly higher in CS₁ and lowest in CS₄. SOC was 19% higher in CS₇ and BD was 16% lower in CS₄ compared to CS₁. Soil porosity and water-stable aggregates (WSA) were also found significantly higher in CS₄ and CS₇ whereas lowest in CS₁. The increase in soil depths significantly increased the soil pH, BD and PR, whereas decreased the soil EC, SOC, AAC, porosity, MWD, and WSA. The cropping systems (CS₄, CS₇, CS₈, CS₆, CS₅, CS₃) with green manure/legume/fodder crops in rotation resulted in better physico-chemical properties through the addition of organic matter to the soil compared to cereal-cereal rotations.

Keywords: *Cropping systems; bulk density; soil organic carbon; mean weight diameter; penetration resistance; green manure.*

1. INTRODUCTION

Rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) cropping system was hitherto a popular practice in northwestern India, especially in Punjab, Haryana, and Uttar Pradesh, because of favourable climatic conditions. This was encouraged by high-yielding rice varieties, traditionally used, and availability of abundant groundwater for irrigation (Ambast, et al., 2016). This rice-wheat cropping system resulted in higher food-grain production, but currently it is believed to be the cause of decline in soil and water resources as well as endangering the sustainability of this system (Chuhan, et al., 2012). Puddling is a land preparation practice that increases water use efficiency and reduces percolation losses in rice (Singh, et al., 2001, Singh, et al., 2011). Continuous intensive puddling practices for a long time form the hardpan in the sub-surface, which adversely affects the growth of other subsequent crops after rice by restricting the root development, nutrient uptake, and the exchange of air with the atmosphere (Singh, et al., 2009, Kumar, et al., 2020). Continuous cereal-cereal crop rotations, removal of crop residues, and other inappropriate tillage practices cause depletion of nutrients and degradation of soil structure (Mamta, et al., 2020, Hiltbrunner, et al., 2013). Cereal-cereal crop rotations are more exhaustive than cereal-legume and cereal-oilseed rotations (Kumar, et al., 2012). The soil

becomes looser and more porous with the addition of a higher amount of biomass (Alam & Salahin, 2013) from the diversity of crops. The quality of organic matter is more important than its quantity for improving the physical condition of the soil (Nweke & Nnabude, 2015). The cropping systems with the inclusion of green manures or legume crops add more soil organic matter as per crop, which further improves soil physical properties (Demir & Isik 2019, Ram, et al., 2022) and C sequestration (Acosta-Martinez, et al., 2011) depending upon the type and quantity of crop residue added to the soil (Zuber, et al., 2015). Significant difference was reported for soil pH (Trehan, et al., 2001) and soil EC values (Kumar, et al., 2020) from various cropping systems. Lowest pH values were reported from the cropping systems adding higher amount of organic matter (Degu, et al., 2019). So, different cropping systems may affect the soil physico-chemical properties differently. The cropping systems add different amounts of organic matter to the soil, on its decomposition tends to variations in soil organic carbon levels (Paranychianakis, et al., 2021). Soil organic matter maintains soil structure, nutrient cycling; acts as a carbon sink, and mitigates the effects of greenhouse gases, thorough specific land use practices (Lal, 2004).

Soil organic carbon (SOC) is an important component in soil that contributes to soil fertility, soil tilth, crop production and soil sustainability.

Soil organic matter (SOM) significantly impacts the soil physico-chemical attributes, viz. soil organic carbon, bulk density (BD), and aggregate stability. Biomass products from cereals and legumes contribute a substantial amount of organic carbon to the soil, improving its physical properties and reducing soil deterioration (Jat, et al., 2013). Organic matter binds the primary soil particles and maintains the aggregate stability, increases water storage, and improves the physical properties of the soil. The use of various crops add a variety of root and shoot biomass, which creates biopores in the soil profile, further reduces the soil compaction and decreases the soil bulk density (Chen & Weil, 2011), and improves soil aggregation (Ram, et al., 2022). Rice grown in rotation with upland crops like maize and mungbean resulted in more SOC content than rice monocropping (Linh, et al., 2016). Also, green manuring crops (legumes and non-legumes) add a higher amount of organic matter to the soil and improve soil physical properties like bulk density, total porosity, soil aggregation, etc. Cropping systems with green manuring of sesbania and green gram improved the organic matter status of soil, which improved the aggregation and reduced the bulk density (Kumar, et al., 2020). Cover crops (CCs) also improve soil physico-chemical properties (Blanco-Canqui, et al., 2011, Cercioglu, et al., 2018). The incorporation of legumes as a cover crop improved the total porosity and aggregate stability, and reduced the soil compaction by adding a higher amount of organic matter (Haruna, et al., 2020). In this context, there is a need to compare the long-term effects of different cropping systems on soil physico-chemical properties. Therefore, the present study was conducted to compare different cropping systems with the inclusion of green manure/legume/ fodder crops to cereals to find their effect on the physico-chemical properties of soil. It may help to promote the rotations of cereal-cereal cropping systems with green manure/fodder/legume crops to sustain the good physical condition of the soil.

2. MATERIALS AND METHODS

2.1 Study Site

The presented study was conducted during *kharif* and *rabi* seasons (2020-2021) in an ongoing long-term experiment at the research farm of the School of Organic Farming, Punjab Agricultural University (PAU), Ludhiana, since 2017. The experiment site was located at 30.90'96" N latitude and 75.78'80" longitude with an altitude of 247 m from the Mean Sea Level representative of Indo-Gangetic alluvial plains with average rainfall of 110-271 mm. The mean minimum temperature in winter varied from 7.2°C in December 2020 to 10.2°C in February 2021 (weather data 2020-21, PAU, Ludhiana). The basis properties of the soil at the site are given in Table 1.

2.2 Treatment Details

The ongoing long-term experiment with ten cropping systems was selected for the present study. The experiment was laid out in a Randomized Complete Block Design (RCBD) with four replications having a plot size of 10.4 x 6.4 m². The treatments included ten cropping systems, viz. CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra. The details about crop variety, fertilizer doses, sowing and harvesting time of the crops in different cropping systems are given in Table 2.

2.3 Soil Sampling, Processing and Analysis

The soil samples were collected from four soil depths (0-7.5, 7.5-15, 15-22.5, and 22.5-30 cm) under each cropping system after harvesting of

Table 1. Basic soil properties of the experimental site (0-30 cm soil depth)

Soil property	Property value
Soil texture	Loamy sand (80-84% sand, 8-20% silt and 8-10% clay)
Soil pH	7.31-7.70
Electrical conductivity (EC)	0.14-0.22 dSm ⁻¹
Soil organic carbon (SOC)	3.9 g kg ⁻¹ (initial)

Table 2. The details of crop varieties, fertilizer doses applied, sowing and harvesting time of the crops in different cropping systems

Crop	Crop variety grown	*Fertilizer dose, N:P₂O₅:K₂O (kg/ha)	Sowing time	Harvesting time
Rice	PR 124	100:30:30	July 15, 2021	October 20, 2021
Wheat	HD 2967	120:60:30	November 13, 2020	April 19, 2021
Basmati rice	Pusa basmati 1121	62.5:0:0	July 27, 2021	November 12, 2021
Cowpea green manure	CL 367	20:50:0	April 27, 2021	June 6, 2021
Mustard	GL 7	120: 60:30	November 11, 2020	March 24, 2021
Maize	PMH 1	120:60:30	July 21, 2021	October 24, 2021
Potato	Pukhraj	188:65:65	October 19, 2020	January 15, 2021
Spring Groundnut	TG 37 A	15:20:25	March 29, 2021	July 23, 2021
Peas	Pb 89	50:65:0	November 10, 2020	March 22, 2021
Oats Fodder	OL 10	30:20:0	November 13, 2020	March 22, 2021
Sathi Maize Fodder	Sathi maize	120:60:30	April 15, 2021	June 21, 2021
Sorghum multicut Fodder	SL 44	100:20:0	April 23, 2020	-
Barseem Fodder	BL 42	25:75:0	September 24, 2020	April 20, 2021
Onion	Punjab Naroya	100:50:50	January 25, 2021	May 25, 2021
Baby corn	PMH 2	60:0:0	July 27, 2021	September 27, 2021
Okra	Punjab Suhawani	90:0:0	February 25, 2021	May 25, 2021

**The nutrient doses were applied as per the recommendations by Punjab Agricultural University, Ludhiana. Urea, diammonium phosphate, and muriate of potash were used as fertilizer sources for N, P₂O₅, and K₂O respectively.*

kharif and *rabi* crops during 2020-21. The collected samples were air-dried, ground, sieved through 2 mm-sized sieves, and stored for analysis. Soil pH and EC were determined using pH and EC meters (Jackson, 1973). Soil organic carbon (SOC) was estimated by the rapid titration method (Walkley & Black, 1934). Undisturbed soil samples were collected with cylindrical iron cores using the core method to determine the soil bulk density (Blake & Hartge, 1986).

Soil porosity was determined by the indirect method (Hillel, 1982) using BD and PD values:

$$\text{Porosity (\%)} = \{(1 - \text{BD}/\text{PD})\} * 100 \quad (1)$$

Where, BD = bulk density

PD = particle density

Aggregate analysis was done using the wet sieving method (Yoder, 1936). Mean weight diameter (Youker & McGuinness, 1957) and water-stable aggregates (Kemper & Rosenau, 1986) were calculated as:

$$\text{MWD} = \sum_{i=1}^n d_i \times w_i \quad (2)$$

$$\text{WSA}_{(0.25 \text{ mm})} = \sum_{i=1}^n n w_i \times 100 \quad (3)$$

Where, n is the number of size fractions, d_i is the mean diameter of each size range (mm), w_i is the weight of aggregates retained in that size range as a fraction of the total dry weight of the sample analyzed (g). Aggregate associated carbon of the aggregates retained on sieve sizes 0.25 mm and 0.1 mm was determined using the rapid titration method (Walkley & Black 1934). Penetration resistance values for all the cropping systems were measured using a hand-held digital cone penetrometer (Naderi-Boldajiet, et al., 2009) after the harvesting of *rabi* and *kharif* crops.

2.4 Statistical Analysis

The data was subjected to statistical analysis by using randomized block design-two-way analysis of variance (ANOVA) in SPSS v 25.0 (SPSS Inc, Chicago, USA). Tukey's honest significant difference test at $p=0.05$ level of significance was used to compare the treatment and depth-wise means.

3. RESULTS AND DISCUSSION

3.1 Soil pH and Soil Electrical Conductivity (EC)

The pooled data pertaining to soil pH in different cropping systems at different soil depths is presented in Figs. 1a and 1b respectively. It depicted that different cropping systems and soil depths significantly ($p=0.05$) affected the soil pH. Comparing all the cropping systems, CS_8 had significantly the lowest pH (7.31) followed by CS_4 (7.37). With the increase in soil depth, it increased significantly in the range of 7.43-7.64. The lowest soil pH in CS_4 and CS_8 is attributed to more decomposition of organic matter added through crop residue compared to other cropping systems. Rice-wheat cropping system (CS_1) had a lower pH than CS_2 , which might be due to submergence in rice as reported by Kumar et al., (2020). The lowest pH in surface soil resulted from N fertilization, which released more H^+ ions during the process of nitrification, in which released nitrate might combined with cations leached from topsoil to subsoil, as these cations were removed and replaced by H^+ ions, and declined the soil pH (Cui, et al., 2022) in surface soil layers. Also, addition of different amount of organic matter by different cropping systems caused the variation in soil pH, as decomposition of organic matter releases several organic acids which are responsible for decrease in soil pH (Ankit, et al., 2024).

The pooled data pertaining to soil EC in different cropping at different soil depths is presented in Figs. 2a and 2b respectively. Soil EC varied in the range 0.14-0.22 dSm^{-1} (Fig. 2a) in different cropping systems and 0.17-0.22 dSm^{-1} at different soil depths (Fig. 2b). Significantly lowest soil EC was observed in CS_7 followed by CS_8 with values of 0.14 and 0.17 respectively. It was significantly higher by 58% in CS_{10} , CS_5 , and by 50% in CS_3 , CS_6 than in CS_7 respectively. It decreased significantly with increase in soil depth. The variations in soil pH might attributed to presence of more crop residues enhances microbial activity, causes anaerobic decomposition of organic matter and produces organic acids which in turn, results subsequent alterations in soil EC (Sharma, et al., 2022). Similar results were also reported by Kumar et al., (2020) that organic matter addition by the different cropping systems affect the salt concentration in the soil and causes variability in soil EC.

3.2 Soil Organic Carbon (SOC) and Mean Weight Diameter (MWD)

The pooled data for SOC and MWD influenced by different cropping systems at different soil depths is given in Table 3. In the last five years (2017-2021), SOC increased from an initial value of 3.9 to 5.0 g kg⁻¹. SOC was significantly (p=0.05) higher in CS₇ (5.45 g kg⁻¹), at par with CS₄ (5.37 g kg⁻¹), followed by CS₆ (5.19 g kg⁻¹) and CS₈ (5.10 g kg⁻¹). The lowest SOC was noticed in CS₁ (4.59 g kg⁻¹) and CS₁₀ (4.65 g kg⁻¹). The highest SOC and MWD in CS₇ resulted from the addition of a higher amount of plant biomass of fodder crops due to higher plant development in both aerial parts and in the roots. Also, a mixture of fodder crops adds more diversified organic residue in the soil which provides favourable conditions for microorganisms and increases carbon accumulation in the soil (Blanco-Canqui & Jasa, 2019, Demir & Isik, 2019). It was also reported by Velso et al., (2020) that the more the presence and diversity of roots, the higher the exudates of organic compounds which serve as a source of soil carbon. In CS₄, there was an additional effect from incorporation of maize crop residue and green manuring of cowpea which added more organic matter to the soil and built up the SOC. Comparing legume-based cropping systems (CS₆, CS₅), the additional effect of two legume crops (peas and spring groundnut) in CS₆ resulted in significantly more SOC than in CS₅. In the cropping systems with potato as a

rabi crop, CS₅ showed more SOC than CS₉ and CS₁₀ due to comparatively more amount of organic matter added from the legume crop in CS₅. Rice-wheat (CS₁) and maize-wheat (CS₂) had comparatively less SOC than CS₃, as over the five years there was no additional source of organic matter to the CS₁ and CS₂, while in CS₃ green manuring was practiced every year before sowing of basmati rice. Depth-wise, SOC decreased significantly from 5.35 to 4.58 g kg⁻¹ at 0-7.5 cm to 22.5-30 cm soil depths respectively. A decrease in SOC with an increase in soil depth was reported (Kumar, et al., 2020, Linh et al., 2016) because of less organic matter addition in lower layers of soil.

MWD was significantly highest in CS₇ with a mean value of 0.342 mm at par with CS₄ (0.329 mm) while lowest in CS₁ (0.182 mm). MWD was higher by 87%, 67%, 65%, and 62% in CS₇ than in CS₁, CS₂, CS₁₀, and CS₉ respectively. Lower MWD in CS₉ and CS₁₀ was caused by intensive tillage in potato crop, which caused the removal of organic matter and decreased soil aggregation.

In rice-based cropping systems, CS₃ (0.205 mm) was at par with CS₁ (0.182 mm), whereas, In the legume-based cropping systems, CS₆ had 29.64% higher MWD than in CS₅. MWD decreased significantly with an increase in soil depth having mean values of 0.330, 0.257, 0.215 and 0.182 mm at 0-7.5, 7.5-15, 15-22.5 and 22.5-30 cm respectively.




Fig. 1 Effect of (a) cropping systems and (b) soil depths on soil pH

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Dissimilar letters indicate the significant difference at p=0.05 by Tukey's honest significant difference




Fig. 2. Effect of (a) different cropping systems and (b) soil depths on soil EC

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Dissimilar letters indicate the significant difference at $p=0.05$ by Tukey's honest significant difference

It was observed that the cropping systems (CS₄, CS₅, CS₆, CS₇, CS₈) with green manure/legume/fodder crops in rotation resulted in higher values for both SOC and MWD compared to other cropping systems. Higher SOC levels increased the MWD in the respective cropping systems as more organic matter addition binds the soil mass together and increases the MWD of soil aggregates. It is also supported by the positive correlation between SOC and MWD (Fig. 7b). Haruna et al., (2020) reported that green manure incorporation in maize crops for 2 years increased the SOC by 0.22%. However, in CS₁, tillage operations for rice puddling destroyed the soil structure and decreased the soil aggregation. Reichert et al., (2022) reported that physical disruption by tillage implements breaks the aggregates, favours more oxidation of organic carbon, and hence reduces soil aggregation. However, fodder crops showed better aggregation, because of more root biomass and aerial parts of the fodder which added organic matter to the soil and provided humic substances as binding agents for aggregate stabilization (Horrocks, et al., 2019).

3.3 Aggregate Associated Carbon (AAC)

The pooled data pertaining to AAC in soil aggregate sizes of 0.25 mm and 0.1 mm in different cropping systems at different soil depths

is presented in Figs. 3a and 3b respectively. In 0.25 mm sized aggregates, AAC was significantly ($p=0.05$) higher in CS₄, which was 11%, 18% and 64% higher than CS₈, CS₇ and CS₁ respectively. The lowest AAC was reported in CS₁ (2.28 mg g⁻¹) followed by CS₁₀ (2.59 mg g⁻¹). Comparing the maize-based cropping systems, the incorporation of green manures/fodder/legume crops in rotation resulted in more AAC in CS₄, CS₆, and CS₇ than in CS₂, and CS₉. In rice-based cropping systems, CS₃ with the incorporation of cowpea as green manure resulted in 39.04% more AAC than CS₁. For aggregate size 0.1 mm, AAC was statistically non-significant ($p=0.05$) in CS₈, CS₄ and CS₇ but significantly higher than CS₁, CS₂, CS₃, CS₉, and CS₁₀ (Fig. 3a). The AAC content in CS₈ was higher by 14%, 16%, and 20% than in CS₉, CS₂, and CS₁₀ respectively. Significantly lowest AAC was reported in CS₁ with a mean value of 2.48 mg g⁻¹ which was lower by 17.9%, 17.3% and 14.5% than in CS₈, CS₄ and CS₇ respectively. AAC in CS₅ was significantly higher by 6.4% and 5.5% than in CS₉ and CS₁₀ attributed to legume crop in CS₅. Saha et al., (2011) revealed that AAC increases with increase in aggregate size, probably because of the larger aggregates formed with the help of organic binding agents from small aggregates. It supported our results that cropping systems with higher organic matter addition resulted more AAC.

Table 3. Effect of different cropping systems on SOC and MWD

Cropping systems	SOC (g kg ⁻¹)				Mean*	MWD (mm)				Mean*
	Soil depths (cm)					Soil depths (cm)				
	0-7.5	7.5-15	15-22.5	22.5-30		0-7.5	7.5-15	15-22.5	22.5-30	
CS ₁	4.89	4.77	4.47	4.24	4.59 ^f	0.226	0.201	0.160	0.143	0.182 ^d
CS ₂	5.17	4.92	4.72	4.44	4.81 ^e	0.277	0.234	0.200	0.176	0.222 ^{cd}
CS ₃	5.22	5.07	4.85	4.54	4.92 ^d	0.247	0.219	0.190	0.164	0.205 ^{cd}
CS ₄	5.77	5.53	5.28	4.88	5.37 ^a	0.472	0.324	0.283	0.238	0.329 ^a
CS ₅	5.27	5.05	4.75	4.54	4.90 ^d	0.317	0.235	0.195	0.160	0.226 ^{cd}
CS ₆	5.61	5.26	5.13	4.78	5.19 ^b	0.427	0.302	0.239	0.205	0.293 ^{ab}
CS ₇	5.88	5.61	5.39	4.93	5.45 ^a	0.492	0.348	0.282	0.247	0.342 ^a
CS ₈	5.59	5.17	4.92	4.72	5.10 ^c	0.350	0.250	0.219	0.176	0.249 ^{bc}
CS ₉	5.08	4.86	4.65	4.39	4.74 ^e	0.251	0.238	0.198	0.158	0.211 ^{cd}
CS ₁₀	5.01	4.79	4.52	4.29	4.65 ^f	0.251	0.221	0.193	0.165	0.207 ^{cd}
Mean*	5.35 ^a	5.10 ^{ab}	4.87 ^{ab}	4.57 ^b		0.331 ^a	0.257 ^b	0.216 ^c	0.183 ^d	

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Means followed by different letters are significantly different at $p=0.05$ by Tukey's honest significant difference




Fig. 3. Effect of (a) cropping systems (b) soil depths on aggregate associated carbon

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Vertical bars and dissimilar letters indicate the significant difference at 5% levels of significance

Depth-wise results for AAC in 0.25 mm and 0.1 mm sized soil aggregates were also significant at given soil depths (Fig. 3b). For both soil aggregate sizes, AAC decreased significantly ($p=0.05$) from surface soil depths to sub-surface. It was higher by 38% and 21% in 0-7.5 cm than in 22.5 cm soil depth for 0.25 mm and 0.1 mm sized soil aggregates respectively.

3.4 Soil Bulk Density (BD)

The pooled data pertaining to soil bulk density in different cropping systems at different soil depths is given in Table 4. Significantly lowest BD was

reported in CS₄ followed by CS₈, CS₇, CS₆ and CS₅ with mean values of 1.41, 1.49, 1.52, 1.54 and 1.55 g cm⁻³ respectively. The highest bulk density was reported in CS₁ (rice-wheat) at par with CS₃ with mean values of 1.66 and 1.64 g cm⁻³ respectively. The lowest BD in CS₄ might result from the soil incorporation of maize residue after harvesting of cobs every year, and the coarse nature of this residue decreased the BD. Additionally to it, cowpea was also incorporated as a green manure from the last five years in CS₄, which added comparatively higher amount of organic matter to the soil. However, higher BD

Table 4. Effect of different cropping systems on soil bulk density

Cropping systems	Soil depths (cm)				Mean*
	0-7.5	7.5-15	15-22.5	22.5-30	
CS ₁ :rice-wheat	1.61	1.65	1.68	1.72	1.66 ^a
CS ₂ : maize-wheat	1.54	1.59	1.63	1.66	1.60 ^{abc}
CS ₃ : basmati rice-wheat-cowpea green manure (GM)	1.57	1.62	1.68	1.70	1.64 ^{ab}
CS ₄ : maize-mustard-cowpea GM	1.32	1.41	1.46	1.48	1.41 ^f
CS ₅ :maize-potato-spring groundnut	1.47	1.53	1.58	1.60	1.55 ^{cde}
CS ₆ : maize-peas-spring groundnut	1.45	1.53	1.59	1.62	1.54 ^{cde}
CS ₇ :maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder	1.42	1.48	1.58	1.62	1.52 ^{de}
CS ₈ : sorghum multicut fodder-barseem fodder	1.37	1.46	1.56	1.57	1.49 ^e
CS ₉ : maize (cobs/fodder)- potato-onion	1.49	1.56	1.61	1.63	1.57 ^{cd}
CS ₁₀ :baby corn-potato-okra	1.52	1.57	1.62	1.65	1.59 ^{bcd}
Mean*	1.47 ^c	1.54 ^b	1.60 ^a	1.62 ^a	

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Means followed by different letters are significantly different at $p=0.05$ by Tukey's honest significant difference

in rice-wheat (CS₁) was caused by puddling in rice carried out every year which might reduce the macroporosity in upper surface layers and resulted in compaction of subsurface layers as also reported by Linh et al., (2016). Intensive tillage practices in potato crop resulted lower BD in CS₉ and CS₁₀ as tillage loosened the soil and resulted less BD than in CS₁. The cropping systems having green manure/fodder/legume crops in rotation resulted in lower BD due to the addition of higher organic matter which added less dense organic components on decaying of roots and crop residues in the soil compared to the mineral constituents of the soil (Beutler, et al., 2017, Santos et al., 2020). BD increased significantly with an increase in soil depths in all the cropping systems. Depth-wise, it varied in the range of 1.47-1.62 g cm⁻³, reported to be lower in surface soil layers compared to sub-surface soil. It was higher by 10% in 22.5 cm soil depth than in 0-7.5 cm due to less organic matter addition in lower layers. The decrease in BD with increased SOC through the addition of organic matter from green manure or fodder crops (Table 3), can be explained by the negative correlation between SOC and BD (Fig. 7a).

3.5 Soil Porosity

The pooled data for soil porosity in different cropping systems at different soil depths is presented in Figs. 4a and 4b respectively. Significantly highest porosity was obtained in

CS₄ at par with CS₈, followed by CS₇ and CS₆ with mean values of 45.10, 42.87, 41.44 and 40.61% respectively. More addition of organic matter from plant biomass increased the soil aggregation, decreased the BD and increased the soil porosity in CS₄ by 5.2%, 8% and 21% compared to CS₈, CS₇ and CS₁ respectively. Rice-wheat (CS₁) cropping system resulted in the lowest porosity with a mean of 37% which was lower by 17%, 13% and 10% than in CS₄, CS₈ and CS₇ respectively. This was due to hardpan formation at the subsurface, which reduced the porosity due to increased compaction in rice-wheat (CS₁) cropping system. Higher porosity in CS₇ and CS₈ is supported by Reichert et al., (2022), who found that fodder crops increased the porosity by improving the proportion of macro aggregates. Soil porosity decreased significantly with an increase in soil depth with mean of 42.95%, 40.79%, 38.91% and 38.31% at 0-7.5, 7.5-15, 15-22.5 and 22.5-30 cm soil depths respectively (Fig. 4b). Porosity was below 40% from depth of 15 - 30 cm, indicating that below this depth plant may not have adequate air and water hence the fields may need deeper tillage to loose the soil and improve the porosity.

3.6 Water Stable Aggregates (0.25 mm) (%)

The pooled data for water-stable aggregates (WSA) in 0.25 mm sized soil aggregates in different cropping systems at different soil depths are shown in Figs. 5a and 5b respectively.




Fig. 4. Effect of (a) cropping systems and (b) soil depths on soil porosity

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Means followed by different letters are significantly different at $p=0.05$ by Tukey's honest significant difference

WSA were reported to be significantly higher ($p=0.05$) in CS₇ at par with CS₄, and CS₈ with mean values of 38.32, 37.42 and 35.11% respectively. However, the lowest WSA was reported in rice-wheat (CS₁), which was lower by 40% than in CS₇. Comparing rice-based cropping systems, WSA increased by 19% in CS₃ compared to CS₁. It was found by Linh et al., (2016) that puddling destroyed the soil aggregation in rice monocropping compared to the rice-mungbean-rice cropping system. Higher SOC levels in CS₇ and CS₄ resulted in more WSA while puddling in rice destructed the soil aggregates and reduced the WSA. Peterson et al., (2002) reported that addition of higher

amount of crop residue/biomass increased the macroaggregation and stability of aggregates. Increase in WSA% in the cropping systems with fodder crops is supported by the findings of Horrocks et al., (2019). WSA decreased significantly with an increase in soil depth with a higher value (39.54%) in 0-7.5 cm soil depth and then decreased to 30.7, 27.29, and 24.22% at 7.5-15, 15-22.5 and 22.5-30 cm soil depths respectively.

3.7 Penetration Resistance (PR)

The data for penetration resistance as affected by different cropping systems is presented in Fig. 6. It revealed that the highest PR was recorded




Fig. 5. Effect of (a) cropping systems and (b) soil depths on water stable aggregates

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra

*Vertical bars and dissimilar letters indicate the significant difference at 5% levels of significance $p=0.05$ by Tukey's honest significant difference

in the rice-wheat (CS₁) cropping system followed by CS₂ and CS₃. However, CS₄ and CS₇ showed lowest PR compared to other cropping systems. At 0-10 cm soil depth, PR varied in the range of 30-350 kPa in different cropping systems. At a subsurface soil depth of 10-20 cm, PR increased to the maximum value in the range of 330 to 512 kPa in CS₄ and CS₁ respectively. Below 20 cm, a steep decrease in PR (512 to 445 kPa) in CS₁, and a slight decrease were reported in CS₂, CS₃ and CS₁₀. However, in CS₄ it increased up to 382

kPa with an increase in soil depth. PR was found to be significantly higher in CS₁ at 15-20 cm soil layer, which might be as a result of hard pan formation due to puddling in rice whereas, in CS₃, this effect was somewhat overcome by cowpea green manuring. CS₄ and CS₇ showed lower penetration resistance which might be due to higher SOC and lower BD levels (Table 3 and 4 respectively). The cropping systems with green manure/legume/fodder crops in rotation showed less penetration resistance than in CS₁ and this




Fig. 6. Effect of different cropping systems on penetration resistance

CS₁: rice-wheat; CS₂: maize-wheat; CS₃: basmati rice-wheat-cowpea green manure (GM); CS₄: maize-mustard-cowpea GM; CS₅: maize-potato-spring groundnut; CS₆: maize-peas-spring groundnut; CS₇: maize+cowpea(fodder)-maize fodder-oats fodder-sathi maize fodder; CS₈: sorghum multicut fodder-barseem fodder; CS₉: maize (cobs/fodder)- potato-onion; CS₁₀: baby corn-potato-okra




Fig. 7. Relationship between (a) SOC and BD, (b) SOC and MWD, (c) MWD and WSA, (d) BD and PR

is expected as organic manure improved soil physical qualities. An increase in BD increased the cohesion of the soil particles, decreased the porosity and increased the risk of soil compaction which led to an increase in PR. The results were also supported by the positive correlation between BD and PR (Fig. 7 d), which showed that penetration resistance increased with an increase in soil bulk density. Cima et al., (2015) reported that rice rotation with other upland crops decreased the bulk density and increased the porosity which further decreased the penetration resistance compared to rice monoculture. It was also reported by Doan et al., (2005) that legume-based cropping systems decreased the soil compaction.

3.8 Correlation between Soil Physico-Chemical Properties

The linear correlation between SOC and MWD, MWD and WSA shows the direct relationship (Figs. 7b, 7c respectively) and indicates that more addition of organic matter increased the SOC content which increased the MWD and WSA. However, SOC and BD were inversely related (Fig. 7a), which indicated that an increase in SOC reduced the BD. BD was directly related to PR (Fig. 7d), indicates that increase in BD increases the PR.

4. CONCLUSION

Rice-wheat cropping system resulted in higher bulk density, high penetration resistance, less SOC, MWD and porosity due to hardpan formation in the sub-surface. However, the inclusion of cowpea as green manure in rotation with basmati rice had more SOC, AAC, and low PR than the rice-wheat cropping system. Also, among maize-based cropping systems, CS₄, CS₇, CS₈, CS₆, and CS₅ resulted in better soil physico-chemical properties compared to CS₂, CS₉ and CS₁₀. The presence of green manures/fodder/legume crops in CS₄, CS₇, CS₈, CS₆, and CS₅ added a higher amount of root and shoot biomass in the soil, increased the SOC, MWD, porosity, WSA, and reduced the bulk density and penetration resistance. In this way, it was concluded that there is the need to include green manure/fodder/legume crops in rotation as these add a higher amount of crop residue or organic matter to the soil and improve the soil physico-chemical properties of the soil compared to cereal-based cropping systems.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Acosta-Martinez, V., Lascano, R., Calderon, F., Jill, D., Booker, T. M., & Zobeck, D. R. (2011). Dryland cropping systems influence the microbial biomass and enzyme activities in a semiarid sandy soil. *Biology and Fertility of Soils*, 47, 655-677. <http://dx.doi.org/10.1007/s00374-011-0565-1>.
- Alam, M. K., & Salahin, N. (2013). Changes in soil physical properties and crop productivity as influenced by different tillage depths and cropping patterns. *Bangladesh Journal of Agricultural Research*, 38, 289-99.
- Ambast, S. K., Tyagi, N. K., & Raul, S. K. (2006). Management of declining groundwater in the Trans Indo-Gangetic Plain (India): some options. *Agricultural Water Management*, 82, 279-296. <https://doi.org/10.1016/j.agwat.2005.06.005>
- Ankit, Prakash, D., Sheoran, S., Yadav, P. K., Kumari, M., Kumar, S., Prajapat, K., Alamri, S., Siddiqui, M.H. & Gupta R. K. (2024). Different cropping systems impact soil health by improving soil biological activities and total organic carbon content, *Archives of Agronomy and Soil Science*, 70:1, 1-24, <http://dx.doi.org/10.1080/03650340.2024.2419035>.
- Beutler, S. J., Pereira, M. G., Tassinari, W. S., Menezes, M. D., Valladares, G. S., & Anjos, L. H. C. (2017). Bulk density prediction for Histosols and soil horizons with high organic matter content. *Revista Brasileira de Ciencia do Solo*, 41, e0160158.

- Blake, G. R., & Hartge, K. H., (1986). Bulk density. *In Methods of Soil Analysis*. pp 363-375. Wisconsin, USA.
- Blanco-Canqui, H., Mikha, M. M., Presley, D. R & Claassen, M. M. (2011). Addition of cover crops enhances no-till potential for improving soil physical properties. *Soil Science Society of America Journal*, 75, 1471-1478.
- Blanco-Canqui, H., & Jara, P. J. (2019). Do grass and legume cover crops improve soil properties in the long term? *Soil Science Society of America Journal*, 83(4), 1181-1187.
<http://dx.doi.org/10.2136/sssaj2019.02.0055>.
- Cercioglu, M., Anderson, S. H., Udawatta, R. P., & Haruna, S. I. (2018). Effects of cover crop and biofuel management on computed topography-measured pore parameters. *Geoderma*, 319, 80-88.
<https://doi.org/10.1016/j.geoderma.2018.01.005>.
- Chauhan, B. S. (2012). Weed ecology and weed management strategies for dry seeded rice in Asia. *Weed Technology*, 26, 1-13.
<https://doi.org/10.1614/WT-D-11-00105.1>.
- Chen, G., & Weil, R. R. (2011). Root growth and yield of maize as affected by soil compaction and cover crops. *Soil Tillage Research*, 117, 17-27.
- Cima, D. S., Luik, A., & Reintam, E. (2015). Organic farming and cover crops as an alternative to mineral fertilizers to improve soil physical properties. *International Agrophysics*, 29, 405-412.
- Cui, H., Luo, Y., Chen, J., Jin, M., Li, Y., & Wang, Z. (2022) Straw return strategies to improve soil properties and crop productivity in a Winter Wheat-Summer Maize cropping system. *European Journal of Agronomy*, 133, 126436.
- Degu, M., Melese, A., & Tena, W. (2019). Effects of soil conservation practice and crop rotation on selected soil physical chemical properties: the case of Dembecha district, Northwestern Ethiopia. *Applied and Environmental Soil Science*, 6, 1-14.
- Demir, Z., & Isik, D. (2019). Effects of cover crops on soil hydraulic properties and yield in a persimmon orchard. *Bragantia*, 78(4), 596-605.
<https://doi.org/10.1590/1678-4499.2010197>.
- Doan, V. Y., Chen, & Irvine, B. (2005). Effect of residue type on the performance of no-till seeder openers. *Canadian Biosystems Engineering*, 472, 2.29-2.35.
- Haruna, S. I., Anderson, S. H., & Udawatta, R. P. (2020). Improving soil physical properties through the use of cover crops: A review. *Agrosystems, Geosciences & Environment* 3, e20105.
<https://doi.org/10.1002/agg2.20105>.
- Hillel, D. (1982). Introduction to Soil Physics. Pp 14 *Academic Press*, New York.
- Hiltbrunner, D., Zimmermann, S., & Hagedorn, F. (2013). Afforestation with Norway spruce on a subalpine pasture alters carbon dynamics but only moderately affects soil carbon storage. *Biochemistry*, 115, 251-266.
- Horrocks, C. A., Arang, J., Arevalo, A., Nunez, J., Cardoso, J. A., & Dungait, J. A. J. (2019). Smart forage selection could significantly improve soil health in the tropics. *Science of the Total Environment*, 688, 609-621.
<http://dx.doi.org/10.1016/j.scitotenv.2019.06.152>.
- Jackson, M. L. (1973). Soil chemical analysis. *New Delhi Prentice Hall*. pp 48-302.
- Jat, M. L., Gathala, M. K., Saharawat, Y. S., Tatarwal, J. P., Gupta, R., & Yadvinder, S. (2013). Double no till and permanent raised beds in maize-wheat rotation of north-western Indo-gangetic plains of India: Effects on crop yields water productivity, profitability and soil physical properties. *Field Crop Research*, 149, 291-299.
- Kemper, W. D. & Rosenau, R. C. (1986). Aggregate stability and size distribution. *Methods of Soil Analysis: Part 1 Physical and Mineralogical Methods*, 5, 425-442.
- Kumar, A., Tripathi, H. P., & Yadav, R. A. (2012). Intensification and diversification in rice (*Oryza sativa*) -wheat (*Triticum aestivum*) cropping system for sustainability. *Indian Journal of Agronomy*, 57, 319-322.
- Kumar, V., Singh, S. K., Singh, P. & Tiwari, S. (2020). Effects of long-term practices of different cropping systems on physico-chemical parameters of soil quality. *International journal of chemical studies*, 81, 136-140.
<https://doi.org/10.22271/chemi.2020.v8.i2r.8921>.

- Lal, R. (2004). Soil Carbon Sequestration Impacts on Global Climate Change and Food Security. *Science*, 304, 1623-1627.
- Linh, T. B., Van, K. L., Elsacker, S. V. & Cornelis, W. M. (2016). Effect of cropping system on physical properties of clay soil under intensive rice cultivation. *Land Degradation and Development* 27, 973-982.
<http://dx.doi.org/10.1002/ldr.2321>.
- Mamta., Kumar, R., Shambhavi, S., Kumar, R., Bairwa, R., Meena, P., & Dahiya, G. (2020). Effect of cropping system and tillage practices on soil physical properties and maize growth. *International journal of chemical studies*, 8, 1372-1378.
<https://doi.org/10.22271/chemi.2020.v8.i2u.8954>.
- Naderi-Boldaji, M., Alimardani, R., Sharifi, A. & Tabatabaeefar, A. (2009). Economical hand-pushed digital cone penetrometer. *International Agrophysics*, 23, 55-60.
- Nweke, I., & Nnabude, P. (2015). Aggregate stability of four soils as evaluated by different indices. *Journal of Experimental Biology and Agricultural Sciences*, 3, 246-252.
- Paranychianakis, N. V. Giannakis, G., Moraetis, D., Tzanakakis, V.A., Nikolaidis, N.P. (2021). Crop litter has a strong effect on soil organic matter sequestration in semi-arid environments. *Sustainability*, 13(23):13278.
<http://dx.doi.org/10.3390/su132313278>.
- Peterson, G. A., Watefall, D. G., Sherrod, L. A., & Shaver, T. M. (2002). Impact of intensive cropping systems on physical properties of surface soils. *Proceedings Great Plains Soil Fertility Symposium*. pp 241-47. University of Nebraska, Lincoln.
- Ram, B., Singh, A. P., Singh, V., Pareek, N., & Gautam, P. (2022). Long term effect of different crop rotations on soil physical properties in a Mollisol. *International journal of phytopharmacy*, 11, 7-11.
<https://doi.org/10.31254/phyto.2022.11102>.
- Reichert, J. M., Corcini, A. L., Awe, G. O., Reinert, D. J., Albuquerque, J. A., Gallarreta, C. C. G., & Docampo, R. (2022). Onion-forage cropping systems on a Vertic Argiudoll in Uruguay: Onion yield and soil organic matter, aggregation, porosity and permeability. *Soil Tillage Research*, 216, 105229.
<https://doi.org/10.1016/j.still.2021.105229>.
- Saha, D., Kukkal, S. S., & Sharma, S. (2011). Landuse impacts on SOC fractions and aggregate stability in Typic Ustochrepts of Northwest India. *Plant Soil* 339: 457-70.
- Santos, O. A. Q., Silva N. E. C., Garcia, A. C., Fagundes, H.S., Diniz, Y. V. F. G., Ferreira, R., & Pereira, M.G. (2020). Impact of land use on Histosols properties in urban agriculture ecosystems of Rio de Janeiro, Brazil. *Revista Brasileira de Ciencia do Solo*, 44, e0200041.
- Sharma, S., Singh, P., Angmo, P., Satpute, S. (2022). Total and labile pools of organic carbon in relation to soil biological properties under contrasting land-use systems in a dry mountainous region. *Carbon Management*, 13(1):352-371.
<http://dx.doi.org/10.1080/17583004.2022.2089236>.
- Singh, B., Humphreys, E., Eberbach, P. L., Katupitiya, A., Singh, Y., & Kukal, S. S. (2011). Growth, yield and water productivity of zero till wheat as affected by rice straw mulch and irrigation schedule. *Field Crops Research*, 121, 209-225.
- Singh, K. B., Gajri, P. R., & Arora, V. K. (2001). Modelling the effects of soil and water management practices on the water balance and performance of rice. *Agricultural water management*, 49, 77-95.
[https://doi.org/10.1016/S0378-3774\(00\)00144-X](https://doi.org/10.1016/S0378-3774(00)00144-X)
- Singh, K. B., Jalota, S. K., & Sharma, B. D. (2009). Effects of continuous rice-wheat rotation on soil properties from four agro-ecosystems of Indian Punjab. *Communications in Soil Science and Plant Analysis*, 40, 2945-2958.
<https://doi.org/10.1080/00103620903221225>
- Trehan, S. P., Sharma, R. C., & Sharma, H. C. (2001). Long term effect of potato based systems, fertilizers and manures on properties of alluvial soil of Punjab. *JIPA*, 28, 207-15.
- Veloso, M. G., Angers, D. A., Chantigny, M. H., & Bayer, C. (2020). Carbon accumulation and aggregation are mediated by fungi in a subtropical soil under conservation agriculture. *Geoderma*, 363, 114-159.
<http://dx.doi.org/10.1016/j.geoderma.2019.114159>.
- Walkley, A. J. & Black, I. A. (1934). An estimation of the degtjareff method for determining soil organic matter and a proposed

- modification of the chromic acid titration method. *Soil Science*, 37, 29-38.
- Yoder, R. E. (1936). A direct method of aggregate analysis of soils and a study of the physical nature of erosion losses 1. *Agronomy Journal*, 28, 337-351.
- Youker, R. E. & McGuinness, J. L. (1957). A short method of obtaining mean weight-diameter values of aggregate analyses of soils. *Soil Science*, 83, 291-294.
- Zuber, S. M., Behnke, G. D., Nafziger, E. D., & Villamil, M. B. (2015). Crop rotation and tillage effects on soil physical and chemical properties in illinois. *Agronomy Journal*, 100, 971-978.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129861>